



Children's Community Nursing Team

Care of a child with cellulitis

Information and guidelines for parents and carers

Cellulitis is an infection in the soft tissue, resulting in redness, swelling pain and discomfort. This fact sheet aims to provide you with information to support you whilst your child is under the care of the Community Team.

It is important that you:

- monitor the cellulitis for any increase in redness, swelling or discomfort
- keep a regular check on his/her temperature with your own thermometer or with the Tempadots provided
- make sure your child has plenty to drink; this may be warm or cold drinks, to stop him/her becoming dehydrated. Offer your child their normal diet but it is usual for them to have a loss of appetite. If your child is reluctant to eat it is important that you encourage him/her to have plenty of drinks
- ensure your child does not play with their IV cannula. Keep it well covered and protected. The nurse will check the IV cannula during each visit. If you do have any problems or concerns refer to the parent and carer guidelines on Home Peripheral Intravenous Therapy and, if necessary, contact the Community Nurse.

Your child may have some pain or discomfort around the area of cellulitis. Your child can be given a dose of Paracetamol or Nurofen for pain relief as directed on the bottle or the packet.

If your child has any of the listed below symptoms you must contact the Community Team immediately

- your child's temperature is above 38c you should give Paracetamol or Nurofen as advised by the nurse or as per instructions on the bottle/packet
- your child's temperature does not come down to below 38c, 45 minutes after Paracetamol or Nurofen has been given
- if your child has an increase in redness, swelling or pain around the area of cellulitis
- if your child is complaining of headaches or feeling unwell
- your child becomes irritable
- your child becomes very pale or mottled with cold hands and feet
- your child starts to vomit
- your child is refusing all drinks and has not passed urine for a period of 4 hours or more during the day

If your child is drowsy or difficult to wake up you must ring for an ambulance by dialling 999 to bring your child to hospital.

Medication

Your child may be prescribed an antibiotic

called _____

The correct dose for your child is _____

This needs to be given every _____ hours for _____ days.

It is important to finish the course of antibiotics.

Your child has been prescribed paracetamol (Calpol/Disprol).

The correct dose for your child is _____

This can be given every _____ hours.

No more than 4 doses should be given in a 24 hour period

Your child has been prescribed Ibuprofen (Nurofen).

The correct dose for your child is _____

This needs to be given every _____ hours.

Your child will remain a Community patient until 24 hours after their IV antibiotic course has been completed. By this time your child should be almost back to normal with either no or only very slight redness or swelling to the area of cellulitis.

If your child is prescribed a course of oral (by mouth) antibiotics it is important they complete the course.

Usually no hospital follow up appointment is necessary unless requested by the doctor.

It would help us if you write down the following information on the chart provided:

- the temperature of your child and the time it was taken
- the time any medication is given (Paracetamol / Nurofen) and how much
- how much your child has eaten or drunk
- if your child has vomited and how much

For further information and advice

Please read the fact sheets on High Temperature and Home Peripheral Intravenous Therapy. Follow all the advice and contact the nurse at any time with any concerns you may have.

Details of how to contact the Community Nurses are in the Children's Community Nursing Team fact sheet.

Please read the patient information leaflets that accompanies prescribed medication for information and risks and side effects.

This fact sheet only gives general information. You must always discuss the individual treatment of your child with the appropriate member of staff. Do not rely on this leaflet alone for information about your child's treatment.

This information can be made available in other languages and formats if requested.

Alder Hey Children's NHS Foundation Trust
Eaton Road
Liverpool
L12 2AP

Tel: 0151 228 4811
www.alderhey.nhs.uk

