

Reference Number: FOI202223/204
From: Private Individual
Date: 27 July 2022
Subject: Numbers of circumcisions and preputioplatic procedures

Q1 Could you please provide the annual numbers of circumcisions and preputioplasties from 2009 to present for boys up to the age of 16?

A1

Year	Number of Preputioplasties in Boys - Up to 16yrs	Number of Circumcisions in Boys - Up to 16yrs
2009	78	320
2010	107	309
2011	173	276
2012	125	257
2013	116	355
2014	107	311
2015	116	291
2016	113	330
2017	188	311
2018	160	234
2019	139	139
2020	81	73
2021	133	127
2022 - up July	68	74
Grand Total	1704	3407

Q2 Additionally, I'd be grateful if you could provide me with any patient information leaflets you distribute to parents/patients on both procedures

A2 Please see attached documents:

- Preputioplasty information sheet 2021
- Preputioplasty Information 2020
- PS02-Circumcision

Preputioplasty for Balanitis Xerotica Obliterans (BXO) in Children

Information sheet for parents

What is this about?

BXO is a scarring condition that affects between 0.01-0.07% of pre-pubertal boys. We do not know why it occurs, but a similar condition also affects the genitalia of ladies.

What does BXO do?

BXO usually causes the foreskin to become thickened and its opening to be narrowed. As a result of this, the foreskin can not be pulled back (retracted). BXO can also affect the 'head' of the penis (the glans) and this can cause the opening that urine passes through (urethral meatus) to become narrowed, so called meatal stenosis. This can cause difficulty in passing urine.

How is it treated?

Mild forms of BXO can be treated by creams and ointments that contain steroids. Your son may have already been treated by this method. When BXO is more severe, it needs an operation to treat it. Traditionally a circumcision was used in all boys who required surgery for BXO. In a circumcision, the foreskin is permanently removed. More recently, a new technique called 'preputioplasty and injection of steroids' has been used. Here, the foreskin is

preserved but the opening is widened to allow it to be pulled back over the glans and an injection of strong steroids is given into the skin to treat the BXO itself.

Which operation is best for my son?

Currently we do not know the answer to this question. Recent studies found that most boys (4 out of 5) treated with preputioplasty and injection of steroids avoided circumcision. However, some boys treated in this way required a repeat of the procedure. It also showed that boys who underwent preputioplasty appeared to have a lower likelihood of needing surgery for meatal stenosis (6%) compared to those who had been circumcised (19%).

What does the operation involve?

The operation of preputioplasty and injection of steroids involves widening the opening of the foreskin by putting 2-3 cuts in the edge and then injecting some strong steroid medicine directly into the skin. The operation is usually performed when your son has had a general anaesthetic, as it would be too uncomfortable to perform with your son awake. After the surgery the foreskin always looks swollen but

the opening is wider and your son should be able to pass urine easily despite the swelling.

Some boys are found to have meatal stenosis at the time of the operation. Your doctor may advise a further operation to widen the meatus at the same time. The doctor will tell you about this if they think that your son may need it.


Are there any 'down-sides'?

There are some risks of the surgery itself that you should consider.

1. There is a small risk of bleeding or infection, this occurs in less than 1 in 50 cases. The risk is lower than for circumcision.
2. After the swelling from the surgery has settled, it is important that the foreskin is pulled back over the glans regularly, we advise to do this at least 3 times per week. If this is not done, there is a higher risk that the foreskin opening will become tight again and your son may need more surgery. This surgery may be a repeat preputioplasty or a circumcision.
3. There is a chance of your son developing meatal stenosis after the surgery. This risk seems to be higher

after a circumcision than after preputioplasty and injection of steroids. About 1 in 10 boys may need surgery to widen the urethral meatus after a preputioplasty (this risk is 1 in 5 after circumcision).

4. The opening of the foreskin will be uneven after the surgery, For most boys this is only seen during foreskin retraction and does not cause any concern. Some boys however find the cosmetic appearance of the foreskin to be unsatisfactory and sometimes they will prefer to have the foreskin removed (circumcision) as a result. This can be done but will mean they need a further operation.



Urology Department

Preputioplasty for Balanitis Xerotica Obliterans (BXO) in Children

What is this leaflet about?

BXO is a scarring condition of the foreskin that affects less than 1 in 100 boys. We do not know why it occurs, but a similar condition can also affect the genitalia of girls.

What does BXO do?

BXO usually causes the foreskin to become thickened and its opening to be narrowed. As a result of this, the foreskin cannot be pulled back (retracted). BXO can also affect the 'head' of the penis (the glans) and this can cause the opening that urine passes through (urethral meatus) to become narrowed, so called meatal stenosis. Narrowing of both the foreskin opening and the urethral meatus can cause difficulty in passing urine.

How is it treated?

Mild forms of BXO can be treated by cream or ointment that contain steroids. Your son may have already been treated by this method. When BXO is more severe, it needs an operation to treat it.

Traditionally a circumcision was used in all boys who required surgery for BXO. In a circumcision, the foreskin is permanently removed. More recently, a technique called 'preputioplasty and injection of steroids' has been used.

Here, the foreskin is preserved but the opening is widened to allow it to be pulled back over the glans and an injection of strong steroid medicine is given into the skin to treat the BXO itself.



Which operation is best for my son?

Currently we do not know the answer to this question.

Recent studies found that most boys (4 out of 5) treated with preputioplasty and injection of steroids have a good result and avoided circumcision.

However, about 1 out of 5 boys treated with preputioplasty and injection of steroids required a repeat procedure or a circumcision within 1-2 years.

The study showed that boys who underwent preputioplasty appeared to have a lower likelihood of needing surgery for meatal stenosis (6%) compared to those who had been circumcised (19%).

What does the operation involve?

The operation of preputioplasty and injection of steroids involves widening the opening of the foreskin by putting 2-3 cuts in the edge and then injecting some strong steroid medicine directly into the skin.

The operation is usually performed when your son has had a general anaesthetic, as it would be too uncomfortable to perform with your son awake.



After the surgery the foreskin always looks very swollen but the opening is wider and your son should be able to pass urine easily despite the swelling. The swelling goes down after a few days.

Some boys are found to have meatal stenosis at the time of the operation. Your doctor may advise a further operation to widen the meatus at the same time. The doctor will tell you about this if they think that your son may need it.

Are there any 'down-sides'?

There are some risks of the surgery itself that you should consider.

1. There is a small risk of bleeding or infection, this occurs in less than 1 in 50 cases. The risk is lower than for circumcision.
2. After the swelling from the surgery has settled, it is important that the foreskin is pulled back over the glans regularly, we advise to do this at least 3 times per week, but ideally every day.

If this is not done, there is a higher risk that the foreskin opening will become tight again and your son may need more surgery. This surgery may be a repeat preputioplasty or a circumcision.

3. There is a chance of your son developing metal stenosis after the surgery. This risk seems to be higher after a circumcision than after preputioplasty and injection of steroids. About 1 in 10 boys may need surgery to widen the urethral meatus after a preputioplasty.
4. The opening of the foreskin will be uneven after the surgery, For most boys this is only seen during foreskin retraction and does not cause any concern.

However, some boys find the cosmetic appearance of the foreskin to be unsatisfactory and sometimes they will chose to have the foreskin removed (circumcision) as a result. This can be done but will mean they need a further operation.

This leaflet only gives general information. You must always discuss the individual treatment of your child with the appropriate member of staff. Do not rely on this leaflet alone for information about your child's treatment.

This information can be made available in other languages and formats if requested.

Alder Hey Children's NHS Foundation Trust
Alder Hey
Eaton Road
Liverpool
L12 2AP

Tel: 0151 228 4811
www.alderhey.nhs.uk

